

Self-adjointness of the twodimensional massless Dirac Hamiltonian and vacuum energy density in the background of a singular magnetic vortex

Yurii A. Sitenko*

Bogolyubov Institute for Theoretical Physics, National Academy of Sciences,
03143 Kyiv, Ukraine

(Ukrainian Journal of Physics, 45, no.4/5, 569-578 (2000))

A massless spinor field is quantized in the background of a singular static magnetic vortex in 2+1-dimensional space-time. The method of self-adjoint extensions is employed to define the most general set of physically acceptable boundary conditions at the location of the vortex. Under these conditions, the vacuum energy density and effective potential in the vortex background are determined.

I. INTRODUCTION

Singular (or contact or zero-range) interaction potentials were introduced in quantum mechanics more than sixty years ago [1–3]. Since that time the attitude of physicists and mathematicians to this subject was varying, starting from "it is impossible", then to "it is evident", and finally arriving at "it is interesting" (for a review see monograph [4]). A mathematically consistent and rigorous treatment of the subject was developed [5], basing on the notion of self-adjoint extension of a Hermitian (symmetric) operator.

Singular interaction is involved in quantum field theory when, for example, a spinor field is quantized in the background of a pointlike magnetic monopole in threedimensional space or a pointlike magnetic vortex in twodimensional space. In these cases the Dirac Hamiltonian,

*Electronic address: yusitenko@bitp.kiev.ua

in contrast to the Schrodinger one, is free from an explicit δ -function singularity; nonetheless the problem of self-adjoint extension of both Dirac and Schrodinger operators arises, albeit for different reasons (see, for example [6]). A distinguishing feature is that a solution to the Dirac equation, unlike that to the Schrodinger one, cannot obey a condition of regularity at the singularity point. It is necessary then to define a boundary condition at this point, and the least restrictive, but still physically acceptable, condition is such that guarantees self-adjointness of the Dirac Hamiltonian. Thus, effects of polarization of the fermionic vacuum in a singular background (such as a pointlike monopole or a pointlike vortex) appear to depend on the choice of the boundary condition at the singularity point, and a set of permissible boundary conditions is labelled, most generally, by the values of self-adjoint extension parameters. In contrast to the Schrodinger case, the extension in the Dirac case does not reflect additional types of interaction but represents complementary information that must be specified when describing the physical attributes of the already posited singular background configuration.

As a consequence, the fermionic vacuum under the influence of a singular background can acquire rather unusual properties: leakage of quantum numbers from the singularity point occurs. While in the case of a monopole there is leakage of charge to the vacuum, which results in the monopole becoming the dyon violating the Dirac quantization condition and CP symmetry [7–11], in the case of a vortex (the Ehrenberg-Siday-Aharonov-Bohm potential [12,13]) the situation is much more complicated, since there is leakage of both charge and other quantum numbers to the vacuum. Apparently, this is due to a nontrivial topology of the base space in the latter case: $\pi_1 = 0$ in the case of a space with a deleted point, and $\pi_1 = \mathbb{Z}$ in the case of a space with a deleted line (or a plane with a deleted point); here π_1 is the first homotopy group and $\mathbb{Z}$ is a set of integer numbers. For a particular choice of the boundary condition at the location of a singular vortex it has been shown that charge [14,15], current [16] and angular momentum [17] are induced in the vacuum. The induced vacuum quantum numbers under general boundary conditions which are compatible with self-adjointness have been considered in Refs. [18–21].

A pointlike static magnetic vortex (the Ehrenberg-Siday-Aharonov-Bohm configuration) in $2 + 1$ dimensional space-time has the form

$$V^1(\mathbf{x}) = -\Phi^{(0)} \frac{x^2}{(x^1)^2 + (x^2)^2}, \quad V^2(\mathbf{x}) = \Phi^{(0)} \frac{x^1}{(x^1)^2 + (x^2)^2}, \quad (1.1)$$

$$\boldsymbol{\partial} \times \mathbf{V}(\mathbf{x}) = 2\pi\Phi^{(0)}\delta(\mathbf{x}), \quad (1.2)$$

where $\Phi^{(0)}$ is the vortex flux in 2π units, i.e. in the London ($2\pi\hbar ce^{-1}$) units, since we use conventional units $\hbar = c = 1$ and coupling constant e is included into vector potential $\mathbf{V}(\mathbf{x})$. The wave function on the plane (x^1, x^2) with punctured singular point $x^1 = x^2 = 0$ obeys the most general condition (see [18] for more details)

$$\langle r, \varphi + 2\pi | = e^{i2\pi\Upsilon} \langle r, \varphi |, \quad (1.3)$$

where $r = \sqrt{(x^1)^2 + (x^2)^2}$ and $\varphi = \arctan(x^2/x^1)$ are the polar coordinates, and Υ is a continuous real parameter which is varied in the range $0 \leq \Upsilon < 1$. It can be shown (see, for example, [17,18]) that Υ as well as $\Phi^{(0)}$ is changed under singular gauge transformations, whereas difference $\Phi^{(0)} - \Upsilon$ remains invariant. Thus, physically sensible quantities are to depend on the gauge invariant combination $\Phi^{(0)} - \Upsilon$ which will be for brevity denoted as the reduced vortex flux in the following.

Thus far the effects of polarization of the massive fermionic vacuum have been studied. In the present paper we find the energy density which is induced by a singular vortex in the massless fermionic vacuum. In the next section, the most general set of physically acceptable boundary conditions at the singularity point $\mathbf{x} = 0$ is defined. In Section III we show how the problem of both ultraviolet and infrared divergences in vacuum characteristics is solved with the help of zeta function regularization. This allows us to get immediately in Section IV the vacuum energy density; also here the effective action and the effective potential are considered. We summarize results and discuss their consequences in Section V. The method of self-adjoint extension is employed to get the boundary condition at the singularity point in Appendix.

II. QUANTIZATION OF A SPINOR FIELD AND THE BOUNDARY CONDITION AT THE LOCATION OF A VORTEX

The operator of the second-quantized spinor field is presented in the form

$$\Psi(\mathbf{x}, t) = \sum_{E_\lambda > 0} e^{-iE_\lambda t} \langle \mathbf{x} | \lambda \rangle a_\lambda + \sum_{E_\lambda < 0} e^{-iE_\lambda t} \langle \mathbf{x} | \lambda \rangle b_\lambda^+, \quad (2.1)$$

where a_λ^+ and a_λ (b_λ^+ and b_λ) are the spinor particle (antiparticle) creation and annihilation operators satisfying anticommutation relations

$$[a_\lambda, a_{\lambda'}^+]_+ = [b_\lambda, b_{\lambda'}^+]_+ = \langle \lambda | \lambda' \rangle, \quad (2.2)$$

and $\langle \mathbf{x} | \lambda \rangle$ is the solution to the stationary Dirac equation

$$H \langle \mathbf{x} | \lambda \rangle = E_\lambda \langle \mathbf{x} | \lambda \rangle, \quad (2.3)$$

H is the Dirac Hamiltonian, λ is the set of parameters (quantum numbers) specifying a state, E_λ is the energy of a state; symbol $\sum$ means the summation over discrete and the integration (with a certain measure) over continuous values of λ . The ground state $|\text{vac}\rangle$ is defined conventionally by equality

$$a_\lambda |\text{vac}\rangle = b_\lambda |\text{vac}\rangle = 0. \quad (2.4)$$

In the case of quantization of a massless spinor field in the background of static vector field $\mathbf{V}(\mathbf{x})$, the Dirac Hamiltonian takes the form

$$H = -i\boldsymbol{\alpha}[\boldsymbol{\partial} - i\mathbf{V}(\mathbf{x})], \quad (2.5)$$

where

$$\boldsymbol{\alpha} = \gamma^0 \boldsymbol{\gamma}, \quad \beta = \gamma^0, \quad (2.6)$$

γ^0 and $\boldsymbol{\gamma}$ are the Dirac γ matrices. In the 2+1-dimensional space-time $(\mathbf{x}, t) = (x^1, x^2, t)$ the Clifford algebra has two inequivalent irreducible representations which can be differed in the following way:

$$i\gamma^0 \gamma^1 \gamma^2 = s, \quad s = \pm 1. \quad (2.7)$$

Choosing the γ^0 matrix in the diagonal form

$$\gamma^0 = \sigma_3, \quad (2.8)$$

one gets

$$\gamma^1 = e^{\frac{i}{2}\sigma_3\chi_s}i\sigma_1e^{-\frac{i}{2}\sigma_3\chi_s}, \quad \gamma^2 = e^{\frac{i}{2}\sigma_3\chi_s}is\sigma_2e^{-\frac{i}{2}\sigma_3\chi_s}, \quad (2.9)$$

where σ_1, σ_2 and σ_3 are the Pauli matrices, and χ_1 and χ_{-1} are the parameters that are varied in the interval $0 \leq \chi_s < 2\pi$ to go over to the equivalent representations.

A solution to the Dirac equation (2.3) with Hamiltonian (2.5) in background (1.1), that obeys condition (1.3), can be presented as

$$\langle \mathbf{x} | E, n \rangle = \begin{pmatrix} f_n(r, E)e^{i(n+\Upsilon)\varphi} \\ g_n(r, E)e^{i(n+\Upsilon+s)\varphi} \end{pmatrix}, \quad n \in \mathbb{Z}, \quad (2.10)$$

where the column of radial functions $\begin{pmatrix} f_n \\ g_n \end{pmatrix}$ satisfies the equation

$$h_n \begin{pmatrix} f_n \\ g_n \end{pmatrix} = E \begin{pmatrix} f_n \\ g_n \end{pmatrix}, \quad (2.11)$$

and

$$h_n = \begin{pmatrix} 0 & e^{i\chi_s}[\partial_r + s(n - \Phi^{(0)} + \Upsilon + s)r^{-1}] \\ e^{-i\chi_s}[-\partial_r + s(n - \Phi^{(0)} + \Upsilon)r^{-1}] & 0 \end{pmatrix} \quad (2.12)$$

is the partial Dirac Hamiltonian. When reduced vortex flux $\Phi^{(0)} - \Upsilon$ is integer, the requirement of square integrability for wave function (2.10) at $r \rightarrow 0$ provides its regularity, rendering the partial Dirac Hamiltonian h_n for every value of n to be essentially self-adjoint. When $\Phi^{(0)} - \Upsilon$ is fractional, the same is valid only for $n \neq n_0$, where

$$n_0 = \llbracket \Phi^{(0)} - \Upsilon \rrbracket + \frac{1}{2} - \frac{1}{2}s, \quad (2.13)$$

$\llbracket u \rrbracket$ is the integer part of a quantity u (i.e., the greatest integer that is less than or equal to u). For $n = n_0$, each of the two linearly independent solutions to Eq.(2.11) meets

the requirement of square integrability at $r \rightarrow 0$. Any particular solution in this case is characterized by at least one (at most both) of the radial functions being divergent as r^{-p} ($p < 1$) at $r \rightarrow 0$. If one of the two linearly independent solutions is chosen to have a regular upper and an irregular lower component, then the other one has a regular lower and an irregular upper component. Therefore, in contrast to operator h_n ($n \neq n_0$), operator h_{n_0} is not essentially self-adjoint ¹. The Weyl - von Neumann theory of self-adjoint operators (see, e.g., Refs. [4,23]) has to be employed in order to consider the possibility of a self-adjoint extension in the case of $n = n_0$. It is shown in Appendix that the self-adjoint extension exists indeed and is parametrized by one continuous real variable denoted in the following by Θ . Thus operator h_{n_0} is defined on the domain of functions obeying the condition

$$\cos(s\frac{\Theta}{2} + \frac{\pi}{4}) \lim_{r \rightarrow 0} (\mu r)^F f_{n_0} = -e^{i\chi s} \sin(s\frac{\Theta}{2} + \frac{\pi}{4}) \lim_{r \rightarrow 0} (\mu r)^{1-F} g_{n_0}, \quad (2.14)$$

where $\mu > 0$ is the parameter of the dimension of inverse length and

$$F = s\{\Phi^{(0)} - \Upsilon\} + \frac{1}{2} - \frac{1}{2}s, \quad (2.15)$$

$\{u\} = u - \llbracket u \rrbracket$ is the fractional part of a quantity u , $0 \leq \{u\} < 1$; note here that Eq.(2.14) implies that $0 < F < 1$, since in the case of $F = \frac{1}{2} - \frac{1}{2}s$ both f_{n_0} and g_{n_0} obey the condition of regularity at $r \rightarrow 0$. Note also that Eq.(2.14) is periodic in Θ with period 2π ; therefore, without a loss of generality, all permissible values of Θ will be restricted in the following to range $-\pi \leq \Theta \leq \pi$.

III. ZETA FUNCTION

In the second-quantized theory the operator of energy is defined as

¹A corollary of the theorem proven in Ref. [22] states that, for the partial Dirac Hamiltonian to be essentially self-adjoint, it is necessary and sufficient that a non-square-integrable (at $r \rightarrow 0$) solution exist.

$$\hat{\mathcal{E}} = \int d^2x \frac{1}{2} [\Psi^+(\mathbf{x}, t), H\Psi(\mathbf{x}, t)]_- = \not\!\!\!\int (E_\lambda a_\lambda^+ a_\lambda - E_\lambda b_\lambda^+ b_\lambda - \frac{1}{2}|E_\lambda|), \quad (3.1)$$

thus the vacuum expectation value of the energy takes the form

$$\mathcal{E} \equiv \langle \text{vac} | \hat{\mathcal{E}} | \text{vac} \rangle = -\frac{1}{2} \not\!\!\!\int |E_\lambda| = -\frac{1}{2} \int d^2x \text{tr} \langle \mathbf{x} | H | \mathbf{x} \rangle. \quad (3.2)$$

The latter expression is ill-defined due to divergences of various kinds. First, there is a bulk divergence resulting from the integration over the infinite twodimensional space. But, even if one considers the vacuum energy density,

$$\mathcal{E}_\mathbf{x} = -\frac{1}{2} \text{tr} \langle \mathbf{x} | H | \mathbf{x} \rangle, \quad (3.3)$$

still it remains to be divergent. There is a divergence at large values of momentum of integration, $k \rightarrow \infty$. To tame this divergence, let us introduce the zeta function density

$$\zeta_\mathbf{x}(z) = \text{tr} \langle \mathbf{x} | H |^{-2z} | \mathbf{x} \rangle, \quad (3.4)$$

which is ultraviolet convergent at sufficiently large values of $\text{Re } z$. However, exactly at these values of $\text{Re } z$ the integral corresponding to Eq.(3.4) is divergent in the infrared region, as $k \rightarrow 0$. To regularize this last divergence, let us introduce fermion mass m , modifying definition (3.4):

$$\zeta_\mathbf{x}(z|m) = \text{tr} \langle \mathbf{x} | \tilde{H} |^{-2z} | \mathbf{x} \rangle, \quad (3.5)$$

where

$$\tilde{H} = -i\boldsymbol{\alpha}[\boldsymbol{\partial} - i\mathbf{V}(\mathbf{x})] + \beta m, \quad (3.6)$$

and it is implied that the complete set of solutions to the equation

$$\tilde{H} \langle \mathbf{x} | \lambda \rangle = \tilde{E}_\lambda \langle \mathbf{x} | \lambda \rangle, \quad (3.7)$$

instead of those to Eq.(2.3), is used.

In the background of a singular magnetic vortex (1.1) – (1.2) the radial functions of the solutions to Eq.(3.7) take the form:

$$\begin{pmatrix} \tilde{f}_n \\ \tilde{g}_n \end{pmatrix} = \begin{pmatrix} \sqrt{1 + m\tilde{E}^{-1}} J_{l-F}(kr) e^{i\chi_s} \\ \text{sgn}(\tilde{E}) \sqrt{1 - m\tilde{E}^{-1}} J_{l+1-F}(kr) \end{pmatrix}, \quad l = s(n - n_0) > 0, \quad (3.8)$$

$$\begin{pmatrix} \tilde{f}_n \\ \tilde{g}_n \end{pmatrix} = \frac{1}{2\sqrt{\pi}} \begin{pmatrix} \sqrt{1 + m\tilde{E}^{-1}} J_{l'+F}(kr) e^{i\chi_s} \\ -\text{sgn}(\tilde{E}) \sqrt{1 - m\tilde{E}^{-1}} J_{l'-1+F}(kr) \end{pmatrix}, \quad l' = s(n_0 - n) > 0, \quad (3.9)$$

$$\begin{pmatrix} \tilde{f}_{n_0}^{(C)} \\ \tilde{g}_{n_0}^{(C)} \end{pmatrix} = \frac{1}{2\sqrt{\pi}[1 + \sin(2\tilde{\nu}_{\tilde{E}}) \cos(F\pi)]} \times \\ \times \begin{pmatrix} \sqrt{1 + m\tilde{E}^{-1}} [\sin(\tilde{\nu}_{\tilde{E}}) J_{-F}(kr) + \cos(\tilde{\nu}_{\tilde{E}}) J_F(kr)] e^{i\chi_s} \\ \text{sgn}(\tilde{E}) \sqrt{1 - m\tilde{E}^{-1}} [\sin(\tilde{\nu}_{\tilde{E}}) J_{1-F}(kr) - \cos(\tilde{\nu}_{\tilde{E}}) J_{-1+F}(kr)] \end{pmatrix}, \quad (3.10)$$

where $k = \sqrt{\tilde{E}^2 - m^2}$, $J_\rho(u)$ is the Bessel function of order ρ and

$$\tan(\tilde{\nu}_{\tilde{E}}) = \text{sgn}(\tilde{E}) \sqrt{\frac{1 - m\tilde{E}^{-1}}{1 + m\tilde{E}^{-1}}} \left(\frac{k}{2\mu} \right)^{2F-1} \frac{\Gamma(1-F)}{\Gamma(F)} \tan \left(s \frac{\Theta}{2} + \frac{\pi}{4} \right), \quad (3.11)$$

$\Gamma(u)$ is the Euler gamma function; note that the radial functions of irregular solution (3.10) satisfy condition (2.14) (see Appendix). Note also that Eqs.(3.8) – (3.10) correspond to the continuum, $|\tilde{E}| > |m|$ ². In addition to them, in the case of

$$\text{sgn}(m) \cos \Theta < 0, \quad (3.12)$$

an irregular solution corresponding to the bound state appears. Its radial functions are

$$\begin{pmatrix} \tilde{f}_{n_0}^{(\text{BS})} \\ \tilde{g}_{n_0}^{(\text{BS})} \end{pmatrix} = \frac{\kappa}{\pi} \sqrt{\frac{\sin(F\pi)}{1 + (2F-1)m^{-1}E_{\text{BS}}}} \begin{pmatrix} \sqrt{1 + m^{-1}E_{\text{BS}}} K_F(\kappa r) e^{i\chi_s} \\ \text{sgn}(m) \sqrt{1 - m^{-1}E_{\text{BS}}} K_{1-F}(\kappa r) \end{pmatrix}, \quad (3.13)$$

where $\kappa = \sqrt{m^2 - E_{\text{BS}}^2}$, $K_\rho(w)$ is the Macdonald function of order ρ and the bound state energy $\tilde{E} = E_{\text{BS}}$ ($|E_{\text{BS}}| < |m|$) is determined implicitly by the equation

$$\frac{(1 + m^{-1}E_{\text{BS}})^{1-F}}{(1 - m^{-1}E_{\text{BS}})^F} = -\text{sgn}(m) \left(\frac{|m|}{2\mu} \right)^{2F-1} \frac{\Gamma(1-F)}{\Gamma(F)} \tan \left(s \frac{\Theta}{2} + \frac{\pi}{4} \right). \quad (3.14)$$

²In a 2+1-, as well as in any odd-, dimensional space-time mass parameter m in Eq.(3.6) can take both positive and negative values.

Regular solutions (3.8) and (3.9) yield the following contribution to zeta function density (3.5):

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{REG}} = \frac{1}{4\pi} \int_0^\infty dk k |\tilde{E}|^{-2z} \sum_{\text{sgn}(\tilde{E})} \left\{ \sum_{l=1}^\infty [(1 + m\tilde{E}^{-1})J_{l-F}^2(kr) + (1 - m\tilde{E}^{-1})J_{l+1-F}^2(kr)] + \sum_{l'=1}^\infty [(1 + m\tilde{E}^{-1})J_{l'+F}^2(kr) + (1 - m\tilde{E}^{-1})J_{l'-1+F}^2(kr)] \right\}. \quad (3.15)$$

Summing over the energy sign and over l and l' , we get the expression

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{REG}} = \frac{1}{\pi} \int_0^\infty dk k |\tilde{E}|^{-2z} \int_0^{kr} \frac{dy}{y} [F J_F^2(y) + (1 - F) J_{1-F}^2(y)], \quad (3.16)$$

which in the case of $\text{Re } z > 1$ is reduced to the form

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{REG}} = \frac{1}{2\pi(z-1)} \int_0^\infty \frac{dk}{k} |\tilde{E}|^{2-2z} [F J_F^2(kr) + (1 - F) J_{1-F}^2(kr)]. \quad (3.17)$$

Irregular solution (4.10) yields the following contribution to Eq.(3.5):

$$\begin{aligned} [\zeta_{\mathbf{x}}(z|m)]_{\text{IRREG}} = & \frac{1}{4\pi} \int_0^\infty dk k |\tilde{E}|^{-1-2z} \{ A\mu^{1-2F} k^{2F} [\tilde{L}_{(+)} - \tilde{L}_{(-)}] J_{-F}^2(kr) + \\ & + A\mu^{1-2F} k^{-2(1-F)} [(m - |\tilde{E}|)^2 \tilde{L}_{(+)} - (m + |\tilde{E}|)^2 \tilde{L}_{(-)}] J_{1-F}^2(kr) + \\ & + 2[(m + |\tilde{E}|) \tilde{L}_{(+)} - (m - |\tilde{E}|) \tilde{L}_{(-)}] J_{-F}(kr) J_F(kr) + 2[(m - |\tilde{E}|) \tilde{L}_{(+)} - \\ & - (m + |\tilde{E}|) \tilde{L}_{(-)}] J_{1-F}(kr) J_{-1+F}(kr) + A^{-1} \mu^{2F-1} k^{-2F} [(m + |\tilde{E}|)^2 \tilde{L}_{(+)} - \\ & - (m - |\tilde{E}|)^2 \tilde{L}_{(-)}] J_F^2(kr) + A^{-1} \mu^{2F-1} k^{2(1-F)} [\tilde{L}_{(+)} - \tilde{L}_{(-)}] J_{-1+F}^2(kr) \}, \end{aligned} \quad (3.18)$$

where summation over the energy sign has been performed and

$$A = 2^{1-2F} \frac{\Gamma(1-F)}{\Gamma(F)} \tan \left(s \frac{\Theta}{2} + \frac{\pi}{4} \right), \quad (3.19)$$

$$\tilde{L}_{(\pm)} = [A\mu^{1-2F} k^{-2(1-F)} (-m \pm |\tilde{E}|) + 2 \cos(F\pi) + A^{-1} \mu^{2F-1} k^{-2F} (m \pm |\tilde{E}|)]^{-1}. \quad (3.20)$$

The contribution of bound state solution (3.13) to Eq.(3.5) is the following:

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{BS}} = \frac{\sin(F\pi)}{\pi^2} \frac{\kappa^2 |E_{\text{BS}}|^{-2z}}{m + E_{\text{BS}}(2F-1)} [(m + E_{\text{BS}}) K_F^2(\kappa r) + (m - E_{\text{BS}}) K_{1-F}^2(\kappa r)]. \quad (3.21)$$

By deforming the contour of integration in the complex w -plane, Eq.(3.17) in the case of $1 < \text{Re } z < 2$ is transformed to the following expression

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{REG}} = \frac{|m|^{2(1-z)}}{2\pi(z-1)} + \frac{\sin(z\pi)}{\pi^2(z-1)} r^{2(z-1)} \times \\ \times \int_{|m|r}^{\infty} \frac{dw}{w} (w^2 - m^2 r^2)^{1-z} [F I_F(w) K_F(w) + (1-F) I_{1-F}(w) K_{1-F}(w)], \quad (3.22)$$

while Eq.(3.18) in the case of $\frac{1}{2} < \text{Re } z < 1$ is transformed to the following one

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{IRREG}} = \frac{\sin(z\pi)}{\pi^2} r^{2(z-1)} \int_{|m|r}^{\infty} dw w (w^2 - m^2 r^2)^{-z} [I_F(w) K_F(w) + I_{1-F}(w) K_{1-F}(w)] + \\ + \frac{2 \sin(F\pi)}{\pi^3} \sin(z\pi) r^{2(z-1)} \times \\ \times \int_{|m|r}^{\infty} dw w (w^2 - m^2 r^2)^{-z} \frac{A \mu^{1-2F} (\frac{w}{r})^{2F} K_F^2(w) + A^{-1} \mu^{2F-1} (\frac{w}{r})^{2(1-F)} K_{1-F}^2(w)}{A \mu^{1-2F} (\frac{w}{r})^{2F} + 2m + A^{-1} \mu^{2F-1} (\frac{w}{r})^{2(1-F)}} - \\ - \frac{\sin(F\pi)}{\pi^2} \frac{\kappa^2 |E_{\text{BS}}|^{-2z}}{m + E_{\text{BS}}(2F-1)} [(m + E_{\text{BS}}) K_F^2(\kappa r) + (m - E_{\text{BS}}) K_{1-F}^2(\kappa r)]; \quad (3.23)$$

here $I_{\rho}(w)$ is the modified Bessel function of order ρ . The integral in Eq.(3.22) can be analytically continued to domain $\frac{1}{2} < \text{Re } z < 2$. In the case of $\frac{1}{2} < \text{Re } z < 1$ this integral is decomposed into two terms:

$$[\zeta_{\mathbf{x}}(z|m)]_{\text{REG}} = \frac{|m|^{2(1-z)}}{2\pi(z-1)} - \\ - \frac{\sin(z\pi)}{\pi^2} r^{2(z-1)} \int_{|m|r}^{\infty} dw w (w^2 - m^2 r^2)^{-z} [I_F(w) K_F(w) + I_{1-F}(w) K_{1-F}(w)] + \\ + \frac{2 \sin(F\pi)}{\pi^3} \frac{\sin(z\pi)}{z-1} r^{2(z-1)} \int_{|m|r}^{\infty} dw (w^2 - m^2 r^2)^{1-z} K_F(w) K_{1-F}(w), \quad (3.24)$$

the last of which can be analytically continued to domain $\text{Re } z < 2$. Note also that the second integral in Eq.(3.23) can be analytically continued to domain $\text{Re } z < 1$.

Summing Eqs.(3.21), (3.23) and (3.24), we get

$$\zeta_{\mathbf{x}}(z|m) = \frac{|m|^{2(1-z)}}{2\pi(z-1)} + \frac{2 \sin(F\pi)}{\pi^3} \frac{\sin(z\pi)}{z-1} r^{2(z-1)} \int_{|m|r}^{\infty} dw (w^2 - m^2 r^2)^{1-z} K_F(w) K_{1-F}(w) +$$

$$\begin{aligned}
& + \frac{2 \sin(F\pi)}{\pi^3} \sin(z\pi) r^{2(z-1)} \times \\
& \times \int_{|m|r}^{\infty} dw w (w^2 - m^2 r^2)^{-z} \frac{A \mu^{1-2F} \left(\frac{w}{r}\right)^{2F} K_F^2(w) + A^{-1} \mu^{2F-1} \left(\frac{w}{r}\right)^{2(1-F)} K_{1-F}^2(w)}{A \mu^{1-2F} \left(\frac{w}{r}\right)^{2F} + 2m + A^{-1} \mu^{2F-1} \left(\frac{w}{r}\right)^{2(1-F)}}, \quad (3.25)
\end{aligned}$$

i.e., the terms which are defined only in domain $\frac{1}{2} < \text{Re } z < 1$ are cancelled.

Note that the first term in Eq.(3.25) is identified with the zeta function density in the noninteracting theory (i.e. in the absence of any boundary condition and any background field):

$$\zeta_{\mathbf{x}}^{(0)}(z|m) = \frac{|m|^{2(1-z)}}{2\pi(z-1)}. \quad (3.26)$$

In the noninteracting theory all vacuum values are simply omitted due to the prescription of normal ordering of the product of operators (see, for example, [24]). Therefore, one has to subtract $\zeta_{\mathbf{x}}^{(0)}$ from $\zeta_{\mathbf{x}}$ for the reasons of consistency. Doing this and removing the infrared regulator mass, we obtain the renormalized zeta function density

$$\begin{aligned}
\zeta_{\mathbf{x}}^{\text{ren}}(z) & \equiv \lim_{m \rightarrow 0} [\zeta_{\mathbf{x}}(z|m) - \zeta_{\mathbf{x}}^{(0)}(z|m)] = \\
& = \frac{\sin(F\pi)}{\pi^3} \sin(z\pi) r^{2(z-1)} \left\{ \frac{\sqrt{\pi}}{4} \frac{\Gamma(1-z)}{\Gamma(\frac{3}{2}-z)} \left[1 - 2 \frac{F(1-F)}{1-z} \right] \Gamma(F-z) \Gamma(1-F-z) + \right. \\
& \quad \left. + \int_0^{\infty} dw w^{1-2z} [K_F^2(w) - K_{1-F}^2(w)] \tanh \left[(2F-1) \ln \left(\frac{w}{\mu r} \right) + \ln A \right] \right\}. \quad (3.27)
\end{aligned}$$

IV. ENERGY DENSITY AND EFFECTIVE POTENTIAL

Recalling the formal expressions for the vacuum energy and zeta function densities, Eqs.(3.3) and (3.4), one can easily deduce that the physical (renormalized) vacuum energy density is expressed through the renormalized zeta function density at $z = -\frac{1}{2}$:

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}} = -\frac{1}{2} \zeta_{\mathbf{x}}^{\text{ren}}\left(-\frac{1}{2}\right). \quad (4.1)$$

In the background of a singular magnetic vortex (1.1) – (1.2), using Eq.(3.27), we get the expression

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}} = \frac{\sin(F\pi)}{2\pi r^3} \left\{ \frac{\frac{1}{2} - F}{6 \cos(F\pi)} \left[\frac{3}{4} - F(1-F) \right] + \right.$$

$$+\frac{1}{\pi^2}\int_0^\infty dw w^2[K_F^2(w)-K_{1-F}^2(w)]\tanh[(2F-1)\ln(\frac{w}{\mu r})+\ln A]\Big\}. \quad (4.2)$$

At noninteger values of reduced vortex flux $\Phi^{(0)} - \Upsilon$ (i.e. at $0 < F < 1$) vacuum energy density (4.2) is positive. At half-integer values of the reduced vortex flux ($F = \frac{1}{2}$) we get

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}}|_{F=\frac{1}{2}} = \frac{1}{24\pi^2 r^3}. \quad (4.3)$$

In the case of $\cos \Theta = 0$ we get

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}} = \frac{\tan(F\pi)}{4\pi r^3}(F - \frac{1}{2}) \left[\frac{1}{3}F(1-F) - \frac{1}{4} \mp \frac{1}{2}(F - \frac{1}{2}) \right], \quad \Theta = \pm s\frac{\pi}{2}. \quad (4.4)$$

If $\cos \Theta \neq 0$, then at large distances from the vortex we get

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}} \underset{r \rightarrow \infty}{=} \frac{\tan(F\pi)}{4\pi r^3}(F - \frac{1}{2}) \left[\frac{1}{3}F(1-F) - \frac{1}{4} + \frac{1}{2}|F - \frac{1}{2}| \right]. \quad (4.5)$$

Going over to imaginary time $t = -i\tau$, let us consider the effective action in 2+1-dimensional Euclidean space-time:

$$\begin{aligned} S_{(2+1)}^{\text{eff}}[\mathbf{V}(\mathbf{x})] &= -\ln\{N^{-1} \int d\Psi d\Psi^+ \exp[-\int d\tau d^2x \Psi^+(-i\beta\partial_\tau - i\beta\tilde{H})\Psi]\} = \\ &= -\ln \text{Det}[(-i\beta\partial_\tau - i\beta\tilde{H})\tilde{m}^{-1}]; \end{aligned} \quad (4.6)$$

here N is a normalization factor, parameter $\tilde{m}$ is inserted just for the dimension reasons, while fermion mass m (see Eq.(3.6)) is introduced in order to tame the infrared divergence.

The real part of the effective action is presented in the form ³

$$\text{Re } S_{(2+1)}^{\text{eff}}[\mathbf{V}(\mathbf{x})] = -\frac{1}{2} \int d\tau d^2x \text{tr} < \mathbf{x}, \tau | \ln[(-\partial_\tau^2 + \tilde{H}^2)\tilde{m}^{-2}] | \mathbf{x}, \tau > . \quad (4.7)$$

Let us define the zeta function density in threedimensional space (x^1, x^2, τ) :

$$\zeta_{\mathbf{x},\tau}(z|m) = \text{tr} < \mathbf{x}, \tau | (-\partial_\tau^2 + \tilde{H}^2)^{-z} | \mathbf{x}, \tau > . \quad (4.8)$$

³The imaginary part of the effective action vanishes in the case of a static background.

Then an ultraviolet regularization of Eq.(4.7) can be achieved by expressing its integrand through Eq.(4.8):

$$-\frac{1}{2}\text{tr} \langle \mathbf{x}, \tau | \ln[(-\partial_\tau^2 + \tilde{H}^2)\tilde{m}^{-2}] | \mathbf{x}, \tau \rangle = \frac{1}{2} \left[\frac{d}{dz} \zeta_{\mathbf{x},\tau}(z|m) \right] \Big|_{z=0} + \frac{1}{2} \zeta_{\mathbf{x},\tau}(0|m) \ln \tilde{m}^2. \quad (4.9)$$

In the background of a singular magnetic vortex (1.1) – (1.2) we get, similarly to Eq.(3.25), the following expression

$$\begin{aligned} \zeta_{\mathbf{x},\tau}(z|m) &= \frac{|m|^{3-2z}}{4\pi^{\frac{3}{2}}} \frac{\Gamma(z - \frac{3}{2})}{\Gamma(z)} - \\ &- \frac{\sin(F\pi)}{\pi^{\frac{7}{2}}} \cos(z\pi) \frac{\Gamma(z - \frac{3}{2})}{\Gamma(z)} r^{2z-3} \int_{|m|r}^{\infty} dw (w^2 - m^2 r^2)^{\frac{3}{2}-z} K_F(w) K_{1-F}(w) - \\ &- \frac{\sin(F\pi)}{\pi^{\frac{7}{2}}} \cos(z\pi) \frac{\Gamma(z - \frac{1}{2})}{\Gamma(z)} r^{2z-3} \times \\ &\times \int_{|m|r}^{\infty} dw w (w^2 - m^2 r^2)^{\frac{1}{2}-z} \frac{A\mu^{1-2F}(\frac{w}{r})^{2F} K_F^2(w) + A^{-1}\mu^{2F-1}(\frac{w}{r})^{2(1-F)} K_{1-F}^2(w)}{A\mu^{1-2F}(\frac{w}{r})^{2F} + 2m + A^{-1}\mu^{2F-1}(\frac{w}{r})^{2(1-F)}}; \end{aligned} \quad (4.10)$$

note that the first term in Eq.(4.10) corresponds to the case of the noninteracting theory:

$$\zeta_{\mathbf{x},\tau}^{(0)}(z|m) = \frac{|m|^{3-2z}}{4\pi^{\frac{3}{2}}} \frac{\Gamma(z - \frac{3}{2})}{\Gamma(z)}. \quad (4.11)$$

Note also relation

$$\zeta_{\mathbf{x},\tau}(0|m) = 0, \quad (4.12)$$

which ensures the independence of the effective action on $\tilde{m}^2$; thus Eq.(4.9) takes the form

$$\begin{aligned} -\frac{1}{2}\text{tr} \langle \mathbf{x}, \tau | \ln(-\partial_\tau^2 + \tilde{H}^2)\tilde{m}^{-2} | \mathbf{x}, \tau \rangle &= \frac{|m|^3}{6\pi} - \frac{2\sin(F\pi)}{3\pi^3 r^3} \int_{|m|r}^{\infty} dw (w^2 - m^2 r^2)^{\frac{3}{2}} K_F(w) K_{1-F}(w) + \\ &+ \frac{\sin(F\pi)}{\pi^3 r^3} \int_{|m|r}^{\infty} dw w (w^2 - m^2 r^2)^{\frac{1}{2}} \frac{A\mu^{1-2F}(\frac{w}{r})^{2F} K_F^2(w) + A^{-1}\mu^{2F-1}(\frac{w}{r})^{2(1-F)} K_{1-F}^2(w)}{A\mu^{1-2F}(\frac{w}{r})^{2F} + 2m + A^{-1}\mu^{2F-1}(\frac{w}{r})^{2(1-F)}}. \end{aligned} \quad (4.13)$$

The effective potential in the massless theory is defined as

$$\mathcal{U}^{\text{eff}}(\mathbf{x}, \tau) = -\frac{1}{2} \lim_{m \rightarrow 0} \text{tr} \langle \mathbf{x}, \tau | \ln \left[\frac{(-\partial_\tau^2 + \tilde{H}^2)\tilde{m}^{-2}}{(-\partial_\tau^2 - \boldsymbol{\partial}^2 + m^2)\tilde{m}^{-2}} \right] | \mathbf{x}, \tau \rangle. \quad (4.14)$$

Defining the renormalized zeta function density

$$\zeta_{\mathbf{x},\tau}^{\text{ren}}(z) = \lim_{m \rightarrow 0} [\zeta_{\mathbf{x},\tau}(z|m) - \zeta_{\mathbf{x},\tau}^{(0)}(z|m)], \quad (4.15)$$

we get, similarly to Eq.(4.9),

$$\mathcal{U}^{\text{eff}}(\mathbf{x}, \tau) = \frac{1}{2} \left[\frac{d}{dz} \zeta_{\mathbf{x},\tau}^{\text{ren}}(z) \right] \Big|_{z=0} + \frac{1}{2} \zeta_{\mathbf{x},\tau}^{\text{ren}}(0) \ln \tilde{m}^2. \quad (4.16)$$

In the background of a singular magnetic vortex (1.1) – (1.2), using Eq.(4.10), we get

$$\begin{aligned} \zeta_{\mathbf{x},\tau}^{\text{ren}}(z) = & -\frac{\sin(F\pi)}{2\pi^{\frac{7}{2}}} \cos(z\pi) \frac{\Gamma(z - \frac{1}{2})}{\Gamma(z)} r^{2z-3} \times \\ & \times \left\{ \frac{\sqrt{\pi}}{4} \frac{\Gamma(\frac{3}{2} - z)}{\Gamma(2 - z)} \left[1 - \frac{4F(1 - F)}{3 - 2z} \right] \Gamma(\frac{1}{2} - z + F) \Gamma(\frac{3}{2} - z - F) + \right. \\ & \left. + \int_0^\infty dw w^{2(1-z)} [K_F^2(w) - K_{1-F}^2(w)] \tanh[(2F - 1) \ln(\frac{w}{\mu r}) + \ln A] \right\}, \end{aligned} \quad (4.17)$$

in particular

$$\zeta_{\mathbf{x},\tau}^{\text{ren}}(0) = 0, \quad (4.18)$$

and, thence, we arrive at the remarkable relation

$$\mathcal{U}^{\text{eff}}(\mathbf{x}, \tau) = \mathcal{E}_{\mathbf{x}}^{\text{ren}}, \quad (4.19)$$

where $\mathcal{E}_{\mathbf{x}}^{\text{ren}}$ is given by Eq.(4.2).

Although the last relation looks rather natural and even evident, let us emphasize here that it is a consequence of the relation between the renormalized zeta function densities of different spatial dimensions,

$$\left[\frac{d}{dz} \zeta_{\mathbf{x},\tau}^{\text{ren}}(z) \right] \Big|_{z=0} = -\zeta_{\mathbf{x}}^{\text{ren}}(-\frac{1}{2}), \quad (4.20)$$

and relation (4.18). As it has been shown in Ref. [25], relation (4.20) can be in general broken in spaces of higher dimensions. Moreover, both left- and right-hand sides of Eq.(4.20) can have nothing to do with the true vacuum energy density. Fortunately, this is not relevant for the case considered in the present paper, and, indeed, in the background of a singular magnetic vortex the vacuum energy density coincides with the effective potential.

V. CONCLUSION

In the present paper we show that the massless fermionic vacuum under the influence of a singular magnetic vortex (1.1) – (1.2) in 2+1-dimensional space-time attains the energy density (4.2) which decreases at large distances from the vortex as inverse power with integer exponent (4.5).

The most general set of boundary conditions at the location of the vortex is used, see Eq.(2.14), providing the self-adjointness of the Dirac Hamiltonian; thus the vacuum energy density is depending on self-adjoint extension parameter Θ or A (3.19). As to the dependence on vortex flux $\Phi^{(0)}$, it has been already anticipated in Introduction that all vacuum polarization effects are gauge invariant and thus depend on reduced vortex flux $\Phi^{(0)} - \Upsilon$ rather than on $\Phi^{(0)}$ or Υ separately. Note also that at half-integer values of $\Phi^{(0)} - \Upsilon$ (i.e. at $F = \frac{1}{2}$) the vacuum energy density is given by Eq.(4.3).

Among the whole variety of boundary conditions which are specified by self-adjoint extension parameter Θ , condition $\cos \Theta = 0$ (or $\Theta = \pm \frac{\pi}{2}$) is distinguished, since it corresponds to one of the two components of a solution to the Dirac equation being regular for all n : if $\Theta = s\frac{\pi}{2}$, then the lower components are regular, and, if $\Theta = -s\frac{\pi}{2}$, then the upper components are regular. This condition is parity invariant, and under it the vacuum energy density is given by Eq.(4.4). It should be noted that this condition is extensively discussed in the literature, being involved into the two most popular ones: the condition of maximal simplicity [26]

$$\Theta = \begin{cases} s\frac{\pi}{2}, & s(\Phi^{(0)} - \Upsilon) > 0 \\ -s\frac{\pi}{2}, & s(\Phi^{(0)} - \Upsilon) < 0 \end{cases} \quad (5.1)$$

and the condition of minimal irregularity [15,18]

$$\Theta = \left\{ \begin{array}{ll} s\frac{\pi}{2}, & -\frac{1}{2} < s(\{\Phi^{(0)} - \Upsilon\} - \frac{1}{2}) < 0 \\ 0, & \{\Phi^{(0)} - \Upsilon\} = \frac{1}{2} \\ -s\frac{\pi}{2}, & 0 < s(\{\Phi^{(0)} - \Upsilon\} - \frac{1}{2}) < \frac{1}{2} \end{array} \right\}; \quad (5.2)$$

here, both in Eqs.(5.1) and (5.2), it is implied that $\{\Phi^{(0)} - \Upsilon\} \neq 0$.

Under condition (5.1) we get

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}} = \left\{ \begin{array}{ll} \frac{\tan(\{\Phi^{(0)} - \Upsilon\}\pi)}{12\pi r^3} \{\Phi^{(0)} - \Upsilon\} (\frac{1}{4} - \{\Phi^{(0)} - \Upsilon\}^2), & \Phi^{(0)} - \Upsilon > 0 \\ \frac{\tan[(1 - \{\Phi^{(0)} - \Upsilon\})\pi]}{12\pi r^3} (1 - \{\Phi^{(0)} - \Upsilon\}) [\frac{1}{4} - (1 - \{\Phi^{(0)} - \Upsilon\})^2], & \Phi^{(0)} - \Upsilon < 0 \end{array} \right\}. \quad (5.3)$$

Under condition (5.2) we get

$$\mathcal{E}_{\mathbf{x}}^{\text{ren}} = \left\{ \begin{array}{ll} \frac{\tan(\{\Phi^{(0)} - \Upsilon\}\pi)}{24\pi r^3} (\{\Phi^{(0)} - \Upsilon\} - \frac{1}{2}) [3|\{\Phi^{(0)} - \Upsilon\} - \frac{1}{2}| - \\ -2(\{\Phi^{(0)} - \Upsilon\} - \frac{1}{2})^2 - 1], & \{\Phi^{(0)} - \Upsilon\} \neq \frac{1}{2} \\ \frac{1}{24\pi^2 r^3}, & \{\Phi^{(0)} - \Upsilon\} = \frac{1}{2} \end{array} \right\}, \quad (5.4)$$

It is clear that Eq.(5.4), in contrast to Eq.(5.3), is periodic in the value of the vortex flux.

As it should be expected, the vacuum energy density is invariant under transitions to equivalent representations of the Clifford algebra (i.e. independent of χ_s). It should be emphasized that the vacuum energy density is also invariant under the transition to an inequivalent representation (i.e. under $s \rightarrow -s$).

ACKNOWLEDGEMENTS

I am thankful to R.Jackiw, H.Leutwyler and W Thirring for interesting discussions. The research was supported by the State Foundation for Fundamental Research of Ukraine.

APPENDIX

Let us consider a general case of massive Hamiltonian $\tilde{H}$ (3.6). The relevant partial Hamiltonian has the form

$$\tilde{h}_{n_0} = \begin{pmatrix} m & e^{i\chi_s}[\partial_r + (1-F)r^{-1}] \\ e^{-i\chi_s}(-\partial_r - Fr^{-1}) & -m \end{pmatrix}. \quad (\text{A.1})$$

Let $\tilde{h}$ be the operator in the form of Eq.(A.1), which acts on the domain of functions $\xi^0(r)$ that are regular at $r = 0$. Then its adjoint $\tilde{h}^\dagger$ which is defined by the relation

$$\int_0^\infty dr r [\tilde{h}^\dagger \xi(r)]^\dagger \xi^0(r) = \int_0^\infty dr r [\xi(r)]^\dagger [\tilde{h} \xi^0(r)] \quad (\text{A.2})$$

acts on the domain of functions $\xi(r)$ that are not necessarily regular at $r = 0$. So the question is, whether the domain of definition of $\tilde{h}$ can be extended, resulting in both the operator and its adjoint being defined on the same domain of functions? To answer this, one has to construct the eigenspaces of $\tilde{h}^\dagger$ with complex eigenvalues. They are spanned by the linearly independent square-integrable solutions corresponding to the pair of purely imaginary eigenvalues,

$$\tilde{h}^\dagger \xi^\pm(r) = \pm i\mu \xi^\pm(r), \quad (\text{A.3})$$

where $\mu > 0$ is inserted for the dimension reasons. It can be shown that in the case of Eq.(A.1) only one pair of such solutions exists, thus the deficiency index of $\tilde{h}$ is equal to (1,1). This pair is given by the following expression

$$\xi^\pm(r) = \frac{1}{N} \begin{pmatrix} e^{i\chi_s} \exp[\pm \frac{i}{2} \text{sgn}(m)\eta] K_F(\tilde{\mu}r) \\ \text{sgn}(m) \exp[\mp \frac{i}{2} \text{sgn}(m)\eta] K_{1-F}(\tilde{\mu}r) \end{pmatrix}, \quad (\text{A.4})$$

where N is a certain normalization factor and

$$\tilde{\mu} = \sqrt{\mu^2 + m^2}, \quad \eta = \arctan\left(\frac{\mu}{|m|}\right). \quad (\text{A.5})$$

Self-adjoint extended operator $\tilde{h}^{\theta_s}$ is defined on the domain of functions of the form

$$\begin{pmatrix} \tilde{f}_{n_0} \\ \tilde{g}_{n_0} \end{pmatrix} = \xi^0 + c(\xi^+ + e^{i\theta_s} \xi^-), \quad (\text{A.6})$$

where c is a complex parameter and θ_s is a real continuous parameter which depends, in general, on the choice between the two inequivalent representations of the Clifford algebra.

Using the asymptotics of the Macdonald function at small values of the variable, we get

$$\begin{pmatrix} \tilde{f}_{n_0} \\ \tilde{g}_{n_0} \end{pmatrix} \underset{r \rightarrow 0}{\sim} \begin{pmatrix} e^{i\chi_s} \cos \left\{ \frac{1}{2}[\theta_s - \text{sgn}(m)\eta] \right\} 2^F \Gamma(F) (\tilde{\mu}r)^{-F} \\ \text{sgn}(m) \cos \left\{ \frac{1}{2}[\theta_s + \text{sgn}(m)\eta] \right\} 2^{1-F} \Gamma(1-F) (\tilde{\mu}r)^{-1+F} \end{pmatrix}, \quad (\text{A.7})$$

or

$$\begin{aligned} & \left\{ \tan \left[\frac{1}{2}\theta_s - \frac{1}{2}\text{sgn}(m)\eta \right] \sin \eta - \text{sgn}(m) \cos \eta \right\} \lim_{r \rightarrow 0} (\tilde{\mu}r)^F \tilde{f}_{n_0} = \\ & = -e^{i\chi_s} 2^{2F-1} \frac{\Gamma(F)}{\Gamma(1-F)} \lim_{r \rightarrow 0} (\tilde{\mu}r)^{1-F} \tilde{g}_{n_0}. \end{aligned} \quad (\text{A.8})$$

Defining new parameter Θ by means of relation

$$\tan \left(s \frac{\Theta}{2} + \frac{\pi}{4} \right) = \left\{ \tan \left[\frac{1}{2}\theta_s - \frac{1}{2}\text{sgn}(m)\eta \right] \sin \eta - \text{sgn}(m) \cos \eta \right\}^{-1} \left(\frac{2\mu}{\tilde{\mu}} \right)^{2F-1} \frac{\Gamma(F)}{\Gamma(1-F)}, \quad (\text{A.9})$$

we get Eq.(2.14).

Certainly, both θ_s and Θ can be regarded as self-adjoint extension parameters which are to specify the boundary condition at $r = 0$. The use of Θ in this aspect may seem to be more preferable just for the convenience reasons, because Eq.(2.14) looks much simpler than Eq.(A.8). In particular, Eq.(2.14), in contrast to Eq.(A.8), is independent of m and remains explicitly invariant under $s \rightarrow -s$ (Θ is independent of s).

In the limit of $m \rightarrow \pm 0$ Eq.(A.9) takes the form

$$\tan \left(s \frac{\Theta}{2} + \frac{\pi}{4} \right) = -\tan \left[\frac{1}{2}\theta_s + \frac{1}{4}\text{sgn}(m)\pi \right] 2^{2F-1} \frac{\Gamma(F)}{\Gamma(1-F)}, \quad (\text{A.10})$$

then we get

$$\theta_s = s\theta, \quad (\text{A.11})$$

where θ is independent of s .

Concluding this appendix, let us note that all characteristics of the massless fermionic vacuum are depending on A (3.19) rather than on Θ itself, and A is expressed through θ in the following way

$$A = -\tan\left[\frac{1}{2}s\theta + \frac{1}{4}\text{sgn}(m)\pi\right]. \quad (\text{A.12})$$

- [1] H.Bethe and R.Peierls, Proc.Roy.Soc.(London) **A148**, 146 (1935).
- [2] L.H.Thomas, Phys.Rev. **47**, 903 (1935).
- [3] E.Fermi, Ricerca Scientifica **7**, 13 (1936).
- [4] S.Albeverio, F.Gesztesy, R.Hoegh-Krohn and H. Holden, *Solvable Models in Quantum Mechanics* (Springer-Verlag, Berlin, 1988).
- [5] F.A.Berezin and L.D.Faddeev, Sov.Math.Dokl. **2**, 372 (1961).
- [6] R.Jackiw, in *M.A.B. Beg Memorial Volume*, edited by A.Ali and P.Hoodbhoy (World Scientific, Singapore, 1991).
- [7] A.S.Goldhaber, Phys.Rev. **D16**, 1815 (1977).
- [8] C.J.Callias, Phys.Rev. **D16**, 3068 (1977).
- [9] E.Witten, Phys.Let. **86B**, 283 (1979).
- [10] B.Grossman, Phys.Rev.Lett. **50**, 464 (1983).
- [11] H.Yamagishi, Phys.Rev. **D27**, 2383; **D28**, 977 (1983).
- [12] W.Ehrenberg and R.Siday, Proc.Phys.Soc.(London) **B62**, 8 (1949).
- [13] Y.Aharonov and D.Bohm, Phys.Rev. **115**, 485 (1959).

- [14] Yu.A.Sitenko, Sov.J.Nucl.Phys. **47**, 184 (1988).
- [15] Yu.A.Sitenko, Nucl.Phys. **B342**, 655 (1990); Phys.Lett. **B253**, 138 (1991).
- [16] P.Gornicki, Ann.Phys.(N.Y.) **202**, 271 (1990).
- [17] Yu.A.Sitenko and D.G.Rakityansky, Ukrainian Journ.Phys. **41**, 329 (1996); Phys.Atom.Nucl. **60**, 247; 258 (1997).
- [18] Yu.A.Sitenko, Phys.Lett. **B387**, 334 (1996)
- [19] Yu.A.Sitenko, Phys.Atom.Nucl. **60**, 2102 (1997); (E) **62**, 1084 (1999).
- [20] Yu.A.Sitenko and D.G.Rakityansky, Phys.Atom.Nucl. **60**, 1497 (1997); Ukrainian Journ.Phys. **43**, 761 (1998).
- [21] Yu.A.Sitenko, Phys.Atom.Nucl. **62**, 1056 (1999).
- [22] J.Weidmann, Math.Z. **119**, 349 (1971).
- [23] N.I.Akhiezer and I.M.Glazman, *Theory of Linear Operators in Hilbert Space* (Pitman, Boston, 1981) V.2.
- [24] N.N.Bogolyubov and D.V.Shirkov, *Introduction to the Theory of Quantized Fields* (Wiley, New York, 1980).
- [25] Yu.A.Sitenko and D.G.Rakityansky, Phys.Atom.Nucl. **61**, 1497 (1998).
- [26] M.G.Alford, J.March-Russel and F.Wilczek, Nucl.Phys. **B328**, 140 (1989)